

Lecture 10 : Trigonometric Substitution

To solve integrals containing the following expressions;

$$\sqrt{a^2 - x^2} \quad \sqrt{x^2 + a^2} \quad \sqrt{x^2 - a^2},$$

it is sometimes useful to make the following substitutions:

Expression	Substitution	Identity
$\sqrt{a^2 - x^2}$	$x = a \sin \theta, \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \quad \text{or} \quad \theta = \sin^{-1} \frac{x}{a}$	$1 - \sin^2 \theta = \cos^2 \theta$
$\sqrt{x^2 + a^2}$	$x = a \tan \theta, \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \quad \text{or} \quad \theta = \tan^{-1} \frac{x}{a}$	$1 + \tan^2 \theta = \sec^2 \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta, \quad 0 \leq \theta < \frac{\pi}{2} \quad \text{or} \quad \pi \leq \theta < \frac{3\pi}{2} \quad \text{or} \quad \theta = \sec^{-1} \frac{x}{a}$	$\sec^2 \theta - 1 = \tan^2 \theta$

Note The calculations here are much easier if you use the substitution in reverse: $x = a \sin \theta$ as opposed to $\theta = \sin^{-1} \frac{x}{a}$.

Integrals involving $\sqrt{a^2 - x^2}$

We make the substitution $x = a \sin \theta, \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}, \quad dx = a \cos \theta d\theta,$

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = a |\cos \theta| = a \cos \theta \quad (\text{since } -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \text{ by choice.})$$

Example

$$\int \frac{x^3}{\sqrt{4 - x^2}} dx$$

Example

$$\int \frac{dx}{x^2\sqrt{9-x^2}}$$

You can use this method to derive what you already know

$$\boxed{\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} + C}$$

Integrals involving $\sqrt{x^2 + a^2}$

We make the substitution $x = a \tan \theta$, $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, $dx = a \sec^2 \theta d\theta$,
 $\sqrt{x^2 + a^2} = \sqrt{a^2 \tan^2 \theta + a^2} = a |\sec \theta| = a \sec \theta$ (since $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ by choice.)

Example

$$\int \frac{dx}{\sqrt{x^2 + 4}}$$

You can also use this substitution to get the familiar

$$\boxed{\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + C.}$$

Completing The Square

Sometimes we can convert an integral to a form where trigonometric substitution can be applied by completing the square.

Example Evaluate

$$\int \frac{dx}{\sqrt{x^2 - 4x + 8}}.$$

Integrals involving $\sqrt{x^2 - a^2}$

We make the substitution $x = a \sec \theta$, $0 \leq \theta < \frac{\pi}{2}$ or $\pi \leq \theta < \frac{3\pi}{2}$, $dx = a \sec \theta \tan \theta d\theta$,

This amounts to saying $\theta = \sec^{-1} \frac{x}{a}$,

$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = a |\tan \theta| = a \tan \theta \quad (\text{since } 0 \leq \theta < \frac{\pi}{2} \text{ or } \pi \leq \theta < \frac{3\pi}{2} \text{ by choice.})$$

Example Evaluate

$$\int \frac{1}{x^2 \sqrt{x^2 - 25}} dx$$

Example Evaluate

$$\int_4^6 \frac{1}{\sqrt{x^2 - 6x + 8}} dx$$

You can also use this substitution to get

$$\int \frac{1}{x \sqrt{x^2 - a^2}} dx = \frac{1}{a} \sec^{-1} \frac{x}{a} + C.$$