

Lecture 32: Calculus with Parametric Curves

Let \mathcal{C} be a parametric curve described by the parametric equations $x = f(t), y = g(t)$. If the function f and g are differentiable and y is also a differentiable function of x , the three derivatives $\frac{dy}{dx}$, $\frac{dy}{dt}$ and $\frac{dx}{dt}$ are related by the chain rule:

$$\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt}.$$

Using this we can obtain the formula to compute $\frac{dy}{dx}$ from $\frac{dx}{dt}$ and $\frac{dy}{dt}$:

$$\boxed{\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \quad \text{if} \quad \frac{dx}{dt} \neq 0}$$

1. The value of $\frac{dy}{dx}$ gives the slope of a tangent to the curve at any given point. This sometimes helps us to draw the graph of the curve.
2. The curve has a **horizontal tangent** when $\frac{dy}{dx} = 0$, and has a **vertical tangent** when $\frac{dy}{dx} = \infty$.



The second derivative $\frac{d^2y}{dx^2}$ can be obtained as well from $\frac{dy}{dx}$ and $\frac{dx}{dt}$. Indeed,

$$\boxed{\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}} \quad \text{if} \quad \frac{dx}{dt} \neq 0}$$

Notice the lack of symmetry, to find $\frac{d^2y}{dx^2}$ we divide by the derivative of $\frac{dx}{dt}$ and we do not use the derivative $\frac{dy}{dt}$.

Example 1 (a) Find an equation of the tangent to the curve

$$x = t^2 - 2t \quad y = t^3 - 3t \quad \text{when} \quad t = -2$$



(b) Find the points on the curve where the tangent is horizontal


(c) Does the curve have a vertical tangent?

(d) Use the second derivative to determine where the graph is concave up and concave down.

Example 2 Consider the curve \mathcal{C} defined by the parametric equations

$$x = t \cos t \quad y = t \sin t \quad -\pi \leq t \leq \pi$$

Find the equations of both tangents to \mathcal{C} at $(0, \frac{\pi}{2})$



Area under a curve

Recall that the area under the curve $y = F(x)$ where $a \leq x \leq b$ and $F(x) > 0$ is given by

$$\int_a^b F(x) dx$$

If this curve can be traced by parametric equations $x = f(t)$ and $y = g(t)$, $\alpha \leq t \leq \beta$ then we can calculate the area under the curve by computing the integral:

$$\int_{\alpha}^{\beta} g(t) f'(t) dt \quad \text{or} \quad \int_{\beta}^{\alpha} g(t) f'(t) dt$$

Example Find the area under the curve

$$x = 2 \cos t \quad y = 3 \sin t \quad 0 \leq t \leq \frac{\pi}{2}$$


Arc Length: Length of a curve

If a curve \mathcal{C} is given by parametric equations $x = f(t)$, $y = g(t)$, $\alpha \leq t \leq \beta$, where the derivatives of f and g are continuous in the interval $\alpha \leq t \leq \beta$ and \mathcal{C} is traversed exactly once as t increases from α to β , then we can compute the length of the curve with the following integral:

$$L = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Example Find the arc length of the spiral defined by

$$x = e^t \cos t \quad y = e^t \sin t \quad 0 \leq t \leq 2\pi$$



Example Find the arc length of the circle defined by

$$x = \cos 2t \quad y = \sin 2t \quad 0 \leq t \leq 2\pi$$

Do you see any problems?